

GEOMORPHOLOGICAL QUANTITATIVE ANALYSIS OF SPERCHIOS RIVER BASIN AREA (CENTRAL GREECE) UTILIZING GEOGRAPHICAL INFORMATION SYSTEMS

Psomiadis E.¹, Migiros G.¹ and Antoniou V.¹

¹ *Agricultural University of Athens, Laboratory of Mineralogy & Geology, Iera Odos 75, 118 55, Athens,
mpsomiadis@gmail.com, bagm@aua.gr, vantoniu@gmail.com*

Abstract

In the present study the analysis of the morphometric quantitative parameters of Sperchios river basin, and specifically of the 8 main sub-basins of the northern and southern part, have been made. The integrated use of a Geographic Information System (GIS) allows a thorough spatial analysis of the data derived from digital terrain spatial models that reveal the geomorphological characteristics of an area. The thorough analysis of the results shows the significant difference of the morphological characteristics of the northern and southern part of the catchment area, due to the impact of the neotectonic activity of the area, which creates this asymmetrical topography.

Key words: *Drainage network, morphometry, tectonics, spatial models.*

Περίληψη

Στην παρούσα μελέτη πραγματοποιήθηκε η ανάλυση των μορφομετρικών ποσοτικών παραμέτρων της λεκάνης του ποταμού Σπερχειού, και συγκεκριμένα των 8 κύριων υπολεκάνων του βόρειου και νότιου τμήματος. Η ολοκληρωμένη χρήση ενός Γεωγραφικού Συστήματος Πληροφοριών (GIS) επιτρέπει μια λεπτομερή χωρική ανάλυση των δεδομένων που προκύπτουν από τα ψηφιακά μοντέλα εδάφους, που αναδεικνύουν τα γεωμορφολογικά χαρακτηριστικά μιας περιοχής. Η διεξοδική ανάλυση των αποτελεσμάτων δείχνει τη σημαντική διαφορά των μορφολογικών χαρακτηριστικών του βόρειου και νότιου τμήματος της λεκάνης απορροής, λόγω της επίδρασης της νεοτεκτονικής, όπου δημιουργεί αυτή την ασύμμετρη τοπογραφία.
Λέξεις κλειδιά: *Υδρογραφικό δίκτυο, μορφομετρία, τεκτονική, χωρικά μοντέλα.*

1. Introduction

The development of the drainage network and the morphological features within a river basin affected and dependent on many, complex and interacting parameters (Deffontaines & Chorowicz, 1991). These factors can be classified as internal, external and complex. Essentially, the complex factors are the combination of internal and external factors. Internal factors derived from the behavior of the bedrock and controlled mainly by lithology and structure. The lithology affect through physical (permeability, increased susceptibility to erosion, hardness, etc.) and chemical properties (diagenesis, dissolution, chemistry of groundwater, etc.) of rocks. The factor of structure arises in various ways, such as the geometry, composition and thickness of rocks, and the switch in tectonic regime of rocks, which can create faults and folds (Schumm 1986). The external factors incorporate climate, vegetation and human activity.

The study area is the basin of river Sperchios in Central Greece, which lies between 38° 44' and 39° 05' N, and 21° 50' and 22° 45' N (Figure 1). In the following sections the statistics and quantitatively characteristics of the 8 main sub-basins, 4 at the south part [Roustianiti (Ro), Inachou (In), Gorgopotamou (Go) and Assopou (As)] and 4 at the north part [Vitiolioti (Vi), Archanorema (Ar), Drimarorema (Dr) and Xerias-Lamias (Xe)] of Sperchios river catchment area, are elaborated (Figure 2). Quantitative measurements of the landscape shape and drainage network characteristics allowed us to compare different landforms and to calculate the geomorphic indices that are useful to identify the landscape evolution of the basin. Geographic Information Systems (GIS) technology, focus on efficiently collecting, maintaining, manipulating and analyzing geomorphic information and a variety of spatial data (Minár et al., 2006). In particular, the core of geomorphological analysis using a GIS is based on the fundamental concept of geomorphology which says that activities of geomorphological processes are explicitly represented in the shape of the georelief. This means that analysis of georelief provides correct information about the origin and development of a landscape as well as relevant data about activities of recent and present-day processes. (Psomiadis et al. 2005 ; Psomiadis, 2010).




Figure 1 - Location of the study area [Digital Elevation Model (DEM) of Sperchios river basin].

2. Data - Methodology

For the present study, digital elevation model (DEM), derived spatial data (such as slope, aspect etc), drainage network and structural geology have been used, originated from 12 topographic and geological maps of 1:50.000 scale. Consequently, all the geospatial data were processed and analyzed through ArcGIS software suite v.9.2). The classification of the drainage network was accomplished using Strachler's classification system Strahler (1957). Using the spatial analysis of the Geographic Information System, morphometric parameters for all sub-basins and their characteristics have been calculated and analyzed. In order to investigate the relation between tectonic and drainage network development, the imprint of both high angle normal faults and small-angle reverse faults (overthrust, thrust) was made. Subsequently, a statistical analysis of the linear elements using Georient software v.8.0 and the creation of frequency (rf) and density (rd) rose diagrams, have been accomplished, both for normal and reverse faults (Figure 4) and additionally for each separately stream order of the drainage network, so as to explore the relation between drainage system and tectonic activity.

3. Study area

3.1. Physical Geographical and Morphological Settings

River Sperchios runs through a valley, which is a graben-like asymmetrical depression. Drainage network throughout the basin is dendritic and drains mountains Tymphristos, Oiti and Othrys.




Figure 2 - The distribution of the 8 sub-basins, 4 in the north and 4 in the south part of Sperchios' river catchment area.






Figure 3 Simplified lithological map of Sperchios river basin, including faults and probable faults of the area.



(α) Normal faults



(β) Reverse faults

Figure 4 - Rose diagrams of frequency (rf) and density (rd) for (a) high angle normal faults [(rf=88° & rd=94° (E-W)]; and (b) small-angle reverse faults (overthrust, thrust) [rf=174° & rd=175° (SSE-NNW)].

In the case of Sperchios basin, separating the basin in two parts based on river's flow direction, in north and south part, significant differences between them can be distinguished. According to the diagram of elevation and elevation areas aggregation curves (Figure 5), the southern part exhibits higher relief than the northern part, while presenting more convex curve from 200 to 1.000m, compared with the north in which the corresponding curve are more concave. This curvature of the curve means that from 200 to 1.000m, the area is increasing, as the elevation increases, while it was expected that should decrease. The cause of this anomaly is mainly due to intense tectonism and particularly at the southern part. Moreover, a large area with elevation less than 100m, due to the elongation of Sperchios river central bed and the spread of alluvial deposits can be depicted. Also, significant differences in terms of relief between the southern and the northern part can be distinguished, using a diagram representing the percentage of each elevation zone (Figure 6).




Figure 5- The relation of elevation and elevation areas aggregation (%).




Figure 6 - The percentage of each elevation zone (%).

In the northern part the percentages show highest values in elevations ranging from 400 to 700m, and then decrease rapidly. In contrast in the southern part the percentages normally increase normally up to approximately 1.000m and then decrease smoothly at higher altitudes. The difference in elevation curve for the two parts (south and north) is due to the fact that the main river bed of Sperchios, is a graben-like asymmetrical depression with a WNW-ESE direction, parallel to the active fault of Atalanti, where the southern part is lifted and the northern is sinking (Mariolakos, 1976).

3.2. Geological Setting

The valley of river Sperchios is a part of a tectonic trough, which is controlled by major NW-SE and E-W trending faults parallel to Atalanti normal fault zone. More specifically, from the fault rose diagrams that have been created, it can be observed that high angle normal faults have a direction E-W, while low angle reverse faults (thrust faults) have a NW-SE direction (Figure 4). The high angle normal faults in the south and central part of the basin have a W-E and NW-SE direction, while in the north part have mainly a NE-SW direction. The fault tectonism took place at the end of the Pliocene and beginning of the Pleistocene periods. The area experienced strong earthquakes, like those which occurred in 426 B.C. and 1894 A.D. and induced big disasters in the broader study area (Mouyiaris, 1988; Pirazzoli et al. 1999).

In the drainage basin of River Sperchios, there are three distinct lithological regions according to Ferrière (1977); these are: (i) the western part that is composed of Paleocene-Eocene flysch of the East Pindus and Parnassos geotectonic zones; it is composed of alternating beds of argillite-siltstone-fine conglomerate and intercalations of shale. There is also a limited occurrence of pelagic and marly limestones of the Vardoussia zone in this area, (ii) the south-eastern portion of the basin is composed of Middle Triassic-Jurassic massive dolomites and limestones of the Pelagonian zone; Upper Cretaceous flysch consisting of coarse sandstone alternating with shale and sandy marl belonging to the zone of Beotia; and Upper Cretaceous thick-bedded limestones and Eocene flysch, composed of sandstone, clay and marl of the Parnassos geotectonic zone, and (iii) in the north and north-eastern part, which is formed by an ophiolitic complex in a shale-chert formation composed of shales, cherts and limestones with ultra-basic and basic igneous rocks-peridotite, dunite, pyroxenite, gabbro, serpentine, diabase, dolerite and metamorphosed green phyllites and schists; belonging to Maliakos (Subpelagonian) zone. In addition, Neogene and Quaternary unconsolidated deposits occupy the central and lower (elevations <500 m) part of the elongated drainage basin of the Sperchios river (Maroukian & Lagios, 1987).

4. Analysis and Results of the selected basins

4.1. Northern Part

4.1.1. Vitoliotis

The Vitoliotis stream sub-basin is located in the northwestern part of Sperchios river basin and has a NW-SE direction (Figure 7a). The basin according to its morphometric characteristics (Table 1) is quite mountainous, susceptible according to its shape to flash flood phenomena. The frequency and density of the drainage network show high values, 3.73 and 9.74 respectively. These values indicate a basin which is formed mostly by impermeable rocks (flysch) and a relatively high relief favoring surface runoff and the creation of a dense drainage network. Finally, the sinuosity index is low while instead, in the main stream branch there is a tendency of elongation. The rose diagrams of faults direction indicate that stream of low Strahler's order have been affected by neotectonic activity. The streams of the 4th, 5th and 6th order streams are influenced mainly by low angle reverse faults or older faults of the area (Table 3).

4.1.2. Archanorema

The Archanorema sub-basin is located in the north and central part of Sperchios river basin and has a NW-SE direction (Figure 7b). This basin according to its morphometric characteristics (Table 1) is a small basin (47.4 km²), semi-mountainous, with low relief. The frequency and density of the drainage network show very high values, 3.81 and 10.21 respectively. These values indicate a basin which is characterized by the presence of semi-permeable and impermeable formations (flysch and ophiolites) leading to a dense drainage network. Finally, the sinuosity index is very low indicating the absence of meander. The rose diagrams of faults direction indicate that the streams of 1st, 2nd and of 4th up to 6th order present a SE-NW direction, following the low angle reverse faults. Oppositely, the 3rd order streams, present a SW-NE direction, following the high angle normal faults, affected by neotectonic activity (Table 3)

4.1.3. Drimarorema

The Drimarorema sub-basin is situated in the north-central part of Sperchios river basin and has a NE-SW direction (Figure 7c). This basin according to its morphometric characteristics (Table 1) is a small basin (26.6 km²), with a very low total relief and mean slope value. The elongation ratio shows a tendency of elongation. The frequency and density of the drainage network represent high values, 3.7 and 8.75 respectively. These values indicate a basin which is characterized by the presence of semi-permeable formations (ophiolites) leading to the creation of a dense drainage network. Finally, the sinuosity index is very low indicating the absence of meander. The rose diagrams of faults direction indicate that the streams of 3rd up to 5th order (main branch) present a SE-NW direction, following the low angle reverse faults. Oppositely, the 1st and 2nd order streams, present a W-E and NE-SW direction, following the high angle normal faults, affected by neotectonic activity of the area (Table 3).

4.1.4. Xerias (Lamias)

The Xerias (Lamias) sub-basin is situated in the northeastern part of Sperchios river basin at the west part of Othrys Mountain and has a N-NW to S-SE direction (Figure 7d). This basin according to its morphometric characteristics (Table 1) is a medium to large basin (109.7 km²), the largest of the northern part. It has a high total relief and the elongation ratio shows a tendency of elongation. The frequency and density of the drainage network represent low values, 2.08 and 4.12 respectively. These values indicate a basin which is characterized by the presence of permeable formations (calcareous) leading to the creation of a sparse drainage network. The low values of frequency, is also due to the low relief and slopes of the basin. Finally, the sinuosity index is low indicating the absence of meander. The rose diagrams of faults direction indicate that the streams of 1st up to 4th order present a E-W direction, following the high angle normal faults affected by neotectonic activity of the area, while the main branch of the stream (5th order) present a NE-SW direction following the low angle reverse faults or older faults of the area (Table 3)

4.2. Southern Part

4.2.1. Roustianitis

The Roustianitis sub-basin is situated in the southwestern part of Sperchios river basin and has a SW to NE direction (Figure 8a). The 85% of this sub-basin lies in elevations higher than 600m, while the 75% belongs to the very mountainous area. This basin according to its characteristics (Table 2) is a medium size basin (53.11 km²). It has high values of mean elevation, total relief and steep slopes. Similar high values of relief ratio and slope index is observed. The length-width ratio appears high, presenting an elongated basin, affected probably also by the neotectonic activity of the area. Regarding the frequency and density of the drainage network, this present low values, 2.33 and 3.86 respectively. Oddly, these values are mismatched with the fact that the basin is formed mainly with flysch which gives usually a dense drainage network. Instead of that the drainage network is sparse, due apparently to the presence of the very high relief and the steep

slopes of the basin. The rose diagrams of faults direction indicate that the streams of 1st and 2nd order present a SE-NW and E-W direction, respectively, following the high angle normal faults. On the contrary, the branches of 3rd, 4th and 5th order present a NE-SW direction following the structure of low angle reverse faults (upthrusts, overthrusts) (Table 3).




Figure 7 - The 4 sub-basins of the northern part (a) Vitoliotis, (b) Archanorema, (c) Drimarrorema and (d) Xerias (Lamias). It is presented the geology and the drainage network.

Table 1 - The morphometric characteristics of the northern part sub-basins.

Parameter	Symbol/ Equation	Vitoliotis	Archanorema	Drimarrorema	Xerias (Lamias)
Area (km ²)	(A)	58.85	47.40	26.62	109.74
Basin Length (km)	(Lb)	10.47	12.22	11.84	17.63
Basin width (km)	(Br=A/Lb)	5.62	3.88	2.18	6.22
Perimeter (km)	(P)	36.28	38.54	30.84	57.63
Maximum Elevation hmax (m)	-	1,281.31	826.45	820.5	1,085.51
Minimum Elevation - hmin (m)	-	215.00	100.00	30.8	12.00
Total relief (m)	H= hmax-hmin	1,066.31	726.45	789.70	1,073.51
Median elevation - hmedian (m)	-	670.00	420.00	285.00	593.14
Mean Elevation - hmean (m)	-	747.41	452.64	456.14	540.00
Basin mean slope (%)	BS = $\Sigma Li d/A$	44.00	32.00	28.00	30.00
Relief ratio (RH) (m/km)	RH=H/(Lb)	101.84	59.45	66.70	60.89
Length-Width index (S)	S=(Lb)/(Br)	1.86	3.15	5.43	2.83
Basin shape index	Rf = A/ Lb^2	0.54	0.32	0.19	0.35
Basin circularity ratio	Rc = $4\pi A/P^2$	0.56	0.40	0.35	0.42
Basin elongation ratio	Re = $1.129(A)^{0.5}/Lb$	1.54	0.64	0.49	0.67
Dr. network density index	Co= $0.282P/(A)^{0.5}$	1.33	1.58	1.69	1.55
Number of streams	(Nu)	573.00	484.00	233.00	452.00
Total stream length (km)	(L)	219.77	180.78	98.48	228.48
Main stream length (km)	Lm	11.56	14.46	12.49	19.30
Density (km/km ²)	D=(L)/(A)	3.73	3.81	3.70	2.08
Frequency (1/km ²)	F=(N)/(A)	9.74	10.21	8.75	4.12
Sinuosity index	C= Lm/Lb	1.10	1.18	1.06	1.10
Slope index (m/km)	Sr=H/Lm	92.24	50.23	63.23	55.62

4.2.2. Inachos

The Inachos sub-basin is situated in the southwestern part of Sperchios river basin, covering a very big part of it and has a SW to NE direction (Figure 8b). The main branch of the river, according to the Strahler's classification, is of the 7th order, which intersects with Sperchios, giving the final 8th order that flows into Maliakos gulf. The 89% of this basin lies in elevations higher than 600m, while the 78.5% belongs to mountainous area. This basin according to its morphometric characteristics (Table 2) is the biggest sub-basin of Sperchios (341.86 km²). It has a very high value of mean elevation and total relief. As far as it concerns the frequency and density of the drainage network present very high values, 3.49 and 8.45 respectively. These values are due to the domination of flysch formation in the basin. The high value of frequency indicates also that the basin is influenced by the tectonic activity of local fault zones. The sinuosity is relatively low indicating the absence of meander. The rose diagrams of faults direction indicate that the streams of 1st order show an E-W following the high angle normal faults. The branches of 3rd and 4th order present a S/SE-N/NW direction following the structure of the low angle reverse faults. The branches of 2nd, 3rd, 6th and 7th order present a NE-SW direction, following the normal faults, but they are mainly affected from the existing geological structure (Table 3).

4.2.3. Gorgopotamos

The Gorgopotamos sub-basin is situated in the southeastern part of Sperchios river basin, originates from the eastern part of mountain Oiti and has a W-SW to E-NE direction (Figure 8c). The 76.3% of this basin lies in elevations higher than 600m, while the 74.6% belongs to the mountainous area. A special feature of this basin is the well known from historical events canyon of Gorgopotamos. The creation of the canyon is due to the steep slopes of the basin and the transition of the geological formations from flysch to carbonate rocks. The presence of the impermeable flysch formation at the upper part of the basin causes a very high surface runoff of rainfall water, while the steep slopes of the basin lead to a very fast water flow. At the transition point that embarks on the existence of the carbonate rocks, which are much more erodible than the flysch formation, the creation of the canyon is started up. The whole process is further enhanced by the Sperchios southern part lifting processes, which along with the existing steep slopes favor the retrogressive erosion creation. This basin according to its morphometric characteristics (Table 2), is a medium size basin (67.62 km²) and it has high total relief and steep slopes. The elongation ratio is rather high, presenting an elongated basin, affected probably and by the neotectonic activity of the area. As regards the frequency and density of the drainage network they have medium to high values, 2.75 and 5.87 respectively. These values indicate a basin which is characterized by the presence of permeable calcareous formations, leading to the creation of a sparse drainage network, except from the north part of the basin that flysch formation is appeared. Finally, the sinuosity index is low indicating the absence of meander. The rose diagrams of faults direction indicate that the streams of 1st up to 4th order present mainly a E-W direction, following the high angle normal faults, as well as the main branch of the river (5th order) presenting a NE-SW direction and seems to be affected either from high angle normal faults and other older faults of the area (Table 3).

4.2.4. Assopos

The Assopos sub-basin is situated in the southeastern part of Sperchios river basin, originates from southeastern part of mountain Oiti and has a W-SW to E-NE direction (Figure 8d). In nowadays, subsequently to the infrastructure and drainage constructions that occurred the last decades in the coastal area of Sperchios basin, Assopos is the only river that supply at the downstream part of Sperchios river. The 73.6% of this basin lies in elevations higher than 600m, while the 58.8% belongs to the very mountainous area. The values of the elongation ratio and the length-width index, present an elongated basin, affected by the neotectonic activity of the area. As far as it concerns, the frequency and density of the drainage network have medium values, 2.35 and 4.64 respectively. These values indicate a basin which is characterized by the presence of permeable

calcareous formations that lead to the creation of a sparse drainage network. The high value of the frequency indicates also the basin is influenced by the tectonic activity of the local fault zones. The sinuosity is relatively low and indicates the absence of meander. The rose diagrams of faults direction indicate that the streams of 1st up to 4th order exhibit an E-W and SE-NW direction, following the high angle normal faults, as well as the main branch of the river (5th order) which presents a NE-SW direction and seems to be affected either from high angle normal faults and other older faults of the area (Table 3).




Figure 8 - The 4 sub-basins of the southern part (a) Roustianitis (b)Inachos, (c) Gorgopotamos and (d)Assopos. It is demonstrated the geology and the drainage network.

Table 2 - The morphometric characteristics of the southern part sub-basins.

Parameter	Sym- bol/Equation	Roustiani- tis	Ina- chos	Gor- gopota- mos	As- sopos
Area (km ²)	(A)	53.11	341.86	67.62	112.97
Basin Length (km)	(Lb)	14.61	29.21	15.06	19.45
Basin width (km)	(Br=A/Lb)	3.64	11.70	4.50	5.81
Perimeter (km)	(P)	39.67	104.58	52.90	58.95
Maximum Elevation hmax (m)	-	1,724.61	2,293.64	2,150.61	1,804.25
Minimum Elevation hmin (m)	-	240.50	90	14.00	8.00
Total relief (m)	H= hmax-hmin	1,484.11	2,203.64	2,136.61	1,796.25
Median elevation - hmedian (m)	-	940.00	962.00	1,320.00	680.00

Mean Elevation - hmean (m)	-	998.70	1,155.36	1,089.70	946.94
Basin mean slope (%)	$BS = \Sigma Li d/A$	0.40	0.42	0.41	0.46
Relief ratio (RH) (m/km)	$RH=H/(Lb)$	101.58	75.44	141.87	92.35
Length-Width index (S)	$S=(Lb)/(Br)$	4.01	2.50	3.35	3.35
Basin shape index	$Rf = A/ Lb^2$	0.25	0.40	0.30	0.30
Basin circularity ratio	$Rc = 4\pi A/P^2$	0.42	0.39	0.30	0.41
Basin elongation ratio	$Re = 1.129(A)^{0.5}/ Lb$	0.21	0.72	0.62	0.62
Dr. network density index	$Co=0.282P/(A)^{0.5}$	1.53	1.60	1.82	1.56
Number of streams	(Nu)	205.00	2,887.00	397.00	524.00
Total stream length (km)	(L)	123.64	1,193.69	185.74	259.91
Main stream length (km)	Lm	15.50	30.40	15.88	21.60
Density (km/km2)	$D=(L)/(A)$	2.33	3.49	2.75	2.30
Frequency (1/km2)	$F=(N)/(A)$	3.86	8.45	5.87	4.64
Sinuosity index	$C= Lm/Lb$	1.06	1.04	1.05	1.11
Slope index (m/km)	$Sr=H/Lm$	95.75	72.49	134.54	83.16

Table 3 – Drainage network orientation of frequency (rf) and density (rd) according to the relevant rose diagrams

Parameter	Vi	Ar	Dr	Xe	Ro	In	Go	As	
	North part				South part				
Fre- quenc y and Densi- ty drain- age orien- tation	Order	rf, rd	rf, rd	rf, rd	rf, rd	rf, rd	rf, rd	rf, rd	rf, rd
	1	E-W	S/SE- N/NW	E-W	E-W	SE-NW	E-W	E-W	SE- NW
	2	E-W	S/SE- N/NW	W/NW- E/SE	E-W	E-W	N/NE- S/SW	E-W	SE- NW
	3	E-W	E/NE- W/SW	N/NE- S/SW	W/N W- E/SE	N/NE- S/SW	N/NE- S/SW	W/NW -E/SE	E-W
	4	N/NE - S/SW	SE-NW	N/NE- S/SW	W/N W- E/SE	W/SW- E/NE	S/SE- N/NW	E-W	E-W
	5	S/SE- N/N W	N/NE- S/SW	N/NE- S/SW	NE- SW	NE-SW	S/SE- N/NW	NE-SW	NE- SW
	6	S/SE- N/NW	SE-NW	-	-	-	N/NE- S/SW	-	-
	7	-	-	-	-	-	N/NE- S/SW	-	-

5. Conclusions

Based on a thorough geospatial analysis of the features in the 8 main sub-basins of Sperchios catchment area, a wealth of useful information on the morphology was extracted. In particular it is clear that the sub-basins at the southern part are clearly longer and have relatively smaller width according to their size, while the basins at the northern part, present higher circularity. Additionally, the relief at the south is steeper, with steeper slopes, higher total relief and mean elevation. These characteristics are due probably to the influence of neotectonic activity of the area and the effects of EW faults, which conduce to the lift process of the southern part and the sinking and widening process of the northern part. The drainage network follows these movements and as

a consequence the branches of the streams in the south are longer and sparse, while in the north are shorter, and dense. Moreover, at the evolution processes of the drainage network, the geological background of the area constitutes a fundamental parameter. In Sperchios basin, the western and southwestern part of the basin are dominated by impermeable rocks (flysch), which form a dense drainage network, while in the south-southeastern part of the basin dominated the permeable calcareous rocks, which form a sparse network.

The frequency and density of the drainage network have mostly moderate to high values, mainly due to geological bedrock and the recent ongoing neotectonic activity of the area. Moreover, the highest values of density are noted at the impermeable and semi-permeable geological formations. According to the frequency and density rose diagrams of the drainage network, can be detected that in the sub-basins of the southern part the main orientation is E-W and less E/SE-W/NW and E/NE-W/SW following the high angle normal faults, which affect and form the whole basin of Sperchios. A small exception can be noticed in the sub-basin of Inachos river, where is affected also from older local faults and shows a NE rotation around of a NE-SW axis. This phenomenon influences the drainage network development. Oppositely, at the northern part the drainage network presents a S/SE-N/NW and less NE-SW orientation, following the low angle reverse faults (upthrust, overthrusts). An exception of that is occurred in the western part, which is affected mainly by the normal faults or from older local faults with an E-W orientation.

6. Acknowledgments

During preparation of the manuscript the State Scholarships Foundation of Greece supported one of the authors (E. Psomiadis).

7. References

- Deffontaines B., Chorowicz J., 1991. Principles of drainage basin analysis from multisource data: Application to the structural analysis of the Zaire basin. In: J. Fourniguet & G. Pierre (Eds), Neotectonics. Tectonophysics, 194, pp. 237-263.
- Mariolakos H., 1976. Thoughts and views on certain issues of Geology and Tectonics of the Peloponnese. *Ann. Geol. Pays Hellen.*, 27, 215-313.
- Maroukian H., Lagios E., 1987. Neotectonic movements in the Sperchios River basin, Central Greece. *Z. Geomorph. N.F., Suppl.-Bd.* 63, 133-140.
- Mentlík P., Jedlička K., Minár J., Barka I., 2006. Geomorphological information system: physical model and options of geomorphological analysis. *Geografie, Sborník České Geografické Společnosti*, p. 15-31.
- Mouyiaris N.K., 1988. Destructive historical earthquakes in N. Euboicos and Maliacos Gulfs- Their significance to the evolution of the area. *Proceedings of the Engineering geology of ancient works, monuments and historical sites-Preservation and Protection*, Vol. 3, pp. 1249-1256, Balkema, Rotterdam.
- Pirazzoli P.A., Stiros S.C., Arnold M., Laborel J., Laborel-Deguen F., 1999. Late Holocene coseismic vertical displacements and tsunami deposits near Kynos, gulf of Euboea, central Greece. *Phys. Chem. Earth (A)*, Vol. 24, No 4, 361-367.
- Psomiadis E., Parcharidis I., Stamatis G., Fomelis M., 2005. Remotely sensing data and thematic mapping for sustainable developing in Sperchios river basin (Central Greece). *Proc. of SPIE 5983, Remote Sensing for Environmental Monitoring, GIS Apps and Geology*.
- Psomiadis E., 2010. Geomorphologic and environmental changes research in the Sperchios river basin utilizing new technologies (Remote Sensing & GIS). Doctorate Thesis elaborated at the Agricultural University of Athens, Lab. of Mineralogy & Geology, p. 394.
- Schumm S.A., 1986. Alluvial river response to active tectonics. In: R.E. Wallace (Eds), *Active Tectonics*. Natl. Acad. Press, Washington, D.C., pp. 80-94.
- Strahler N.A., 1957. Quantitative analysis of watershed geomorphology. *Amer. Geophys. Union, Trans.*, 38 (6), 1117-1142.