

succession with lower part of the Ujak succession is covered by thick slope deposits, up to 27 m. Recently, in this area three boreholes have been drilled, by the Polish Geological Institute.

Combining the field observation with core material, following sequence of deposits of the Magura and Ujak transitional zone can be revealed:

1) thick-bedded sandstones of the Piwniczna Member (Early/Middle Eocene) of Magura Formation,

2) thin-bedded flysch and red shales with *Reticulophragmium amplexens* (Mniszek Shale Member, Middle Eocene) of the Magura Formation,

3) few meters thick packet of grey-greenish and red marstone of the Sub-Menilite *Globigerina* Marls (Late Eocene-Early Oligocene),

4) at least, a 19 m thick dark brown and black Menilite Shales with horstone and tuffite intercalations (Oligocene),

5) a 25 m packet of thick-bedded muscovitic sandstones, an equivalent of the Poprad Member (oligocene) of Magura Formation,

6) dark-grey marly shales with intercalations of thin bedded calcareous sandstones of the Malcov Formation (Oligocene).

The studies conducted by us have shown that during the Late Eocene through the Oligocene the Klippen Pieniny Belt was a transitional zone between the Magura and the Central Carpathian basins.

Jurassic and Cretaceous tectonic evolution of the Sakar and Srednogorie zones, Bulgaria: $^{40}\text{Ar}/^{39}\text{Ar}$ mineral ages and structures

Neubauer F.¹, Bilyarski S.², Genser J.¹, Ivanov Z.², Peytcheva I.^{3,4} and von Quadt A.⁴

¹*Department of Geography and Geology, University of Salzburg, A-5020 Salzburg, Austria, franz.neubauer@sbg.ac.at*

²*Dept. of Geology, Sofia University, Sofia, Tzar Osvoboditel Blv. 15, Sofia 1000, Bulgaria*

³*Geological Institute, Bulgarian Academy of Sciences, Acad. G. Bonchev st., 1113 Sofia, Bulgaria*

⁴*Institute of Isotope Geology and Mineral Resources, ETH Zurich, Clausiusstrasse 25, CH-8092 Zurich, Switzerland*

We present new $^{40}\text{Ar}/^{39}\text{Ar}$ mineral ages from metamorphic, plutonic and volcanic rocks of the principally E-W striking the Sakar and Srednogorie zones, Bulgaria and use these ages together with structural observations to constrain the Jurassic and Cretaceous tectonic history of these zones along northern margins of the Rhodope massif. The Srednogorie zone seems to be particularly important because of its richness of magmatic rocks and associated ore deposits.

The Sakar zone SW of Topolovgrad exposes the Palaeozoic Sakar granite mantled by the metamorphic Laslovo Formation. Along the contacts between granite and their volcanosedimentary metamorphic cover, often shear zones developed with a mylonitic fabrics formed within upper greenschist to amphibolite facies-grade metamorphic conditions giving the structure an appearance similar to a mantled gneiss dome. $^{40}\text{Ar}/^{39}\text{Ar}$ amphibole and white mica dating yield ages ranging between 144 and 136 Ma constraining the age of the main tectonic event of ductile deformation within a deep crustal level at ca. the Jurassic/Cretaceous boundary. Further to the southeast, towards the Rhodope massive, younger white mica ages gradually decreasing to ca. 124 Ma were found in metamorphic rocks. A further, secondary thermal overprint was found in staircase Ar release patterns with a maximum age of a secondary thermal overprint younger than 69 Ma. Together, these ages indicate the principal age of the Sakar zone at the Jurassic/Cretaceous boundary, which predates the formation of the Srednogorie basin, and two stages of thermal overprint.

The Srednogorie zone comprises an Upper Cretaceous siliclastic marine infill of a volcanosedimentary basin and abundant volcanics, subvolcanics and shallow plutons. In the eastern Srednogorie zone, alkalic rocks are abundant. A new $^{40}\text{Ar}/^{39}\text{Ar}$ amphibole age of 82 Ma from a hornblende andesite from the Fakjijska river S of Sredec indicate an important stage of effusive volcanic activity. In contrast, the following ages from plutonic rocks are interpreted to date cooling through the Ar retention temperature, ca. 500–550°C for

amphibole and ca. 300°C for biotite after magma crystallization. They represent, therefore, the minimum ages in respect to their crystallization. These ages include: biotite ages of 83, 79 and 79 Ma from the Granitovo granodiorite, Izgrev diorite and Rosen syenite (at Černomorez), respectively; and amphibole ages of ca. 86, 85 and 76 Ma from microgabbro within the Granitovo granodiorite, the Oman-Fakia gabbro (near Danica) and a gabbrodiorite of ENE Samokov. Together, these ages proof the wide range of magmatism between 86 and 76 Ma and a particularly important cooling event in the Srednogorie zone at ca. 80 Ma. In the case of the Izgrev diorite in the eastern Srednogorie zone, cooling at ca. 79 Ma is associated with chalcopyrite mineralization along conjugate shear-extension veins. Their structural assessment proofs NW–SE extension, and this event is in agreement with many similar but poorly dated observations in the eastern Srednogorie zone. This indicates that regional extension was of transtensive type at ca. 80 Ma in respect to the strike of the Srednogorie zone. Inversion of the Srednogorie basin comprises regional N-S shortening and a subsequent stage of NE-SW transpressive motion as abundant sets of conjugate strike-slip faults indicate.

Monitoring the thermal anomalies around Milos Island with satellite thermal data

Nikolakopoulos K.

Division of General Geology and Geological Mapping, Institute of Geology and Mineral Exploration, 1 Sp. Louis Str., Olympic Village Entrance C, 13677 Acharnae, Athens, Greece, knikolakopoulos@igme.gr

Although remote sensing is recognized as a powerful tool in the collection, analysis and modelling of environmental data, less attention has been given to the use of Thermal Infrared (TIR) remote sensing. Thermal property of a material is representative of upper several centimetres of the surface. As in thermal remote sensing we measure the emitted radiations, it proves to be complementary to other remote sensing data and even unique in helping to identify surface materials and features such as rock types, soil moisture, underwater springs, geothermal anomalies etc. During the last two decades a series of satellite and airborne sensors have been developed to collect TIR data from the earth surface, such as HCMM, Landsat TM/ETM+, AVHRR, MODIS, ASTER, and TIMS. In addition to Land Surface Temperature (LST) measurements, these TIR sensors may also be utilized to obtain emissivity data of different surfaces with varied resolutions and accuracies.

The islands of Nisyros, Yali, Kos, Santorini, Milos, Poros, Aegina and the peninsula of Methana constitute the Hellenic Volcanic Island Arc. This arc seems to be geodynamically very active since it comprises the largest volumes of volcanic materials and is at present a region of high tectonic activity. This activity is very often expressed with earthquakes, gas explosions and hydrothermal eruptions, volcanic eruptions, landslides, etc.

TIR data were used in order to detect undersea hydrothermal activities along the Hellenic Volcanic Island Arc. In this study there is a effort to monitor the thermal anomalies into the sea around Milos Island for the last 25 years using TIR data. More especially thermal data from the LANDSAT-TM the LANDSAT-ETM and ASTER sensors were used. The resolution of the thermal infrared bands ranges between 60 and 120 m. The sensitivity of these sensors is about 0.5°C in the region of 10.4-12.5µm (thermal infrared zone) of the Electromagnetic Spectrum. Thus the sensors can contribute to the detection of thermal anomalies (water outflows into the sea environment), which are useful to hydrothermal studies.

All the satellite data were orthorectified. Then using the appropriate algorithms the radiation was converted into Celsius Degrees in order to calculate the surface temperature of the area. Then in order to better distinguish thermal deviations the temperatures have been classified using the density slicing method. The results are presented in this study.